arXiv:0810.0318v2 [math.GM] 3 Jan 2009

On boundedness, existence and uniqueness of strong solutions of the Navier-Stokes Equations in 3 dimenions.

A. A. Ruzmaikina.
Department of Statistics, Purdue University, West Lafayette, IN 47909.
`aar@purdue.edu`

Abstract.

In this paper we consider the Navier-Stokes Equations in $\mathbf{R}^3$ *in the vorticity formulation in the absence of the external forces. We first derive the upper bounds on* $|\omega|_{\frac{6.02}{3}}(t)$ *for* $t \in [t_1, t_2]$ *in terms of* $\max_{t\in[t_1,t_2]}|\omega|_2(t)$*, then derive the upper bounds on* $|\bar\nabla\bar\omega|_2(t)$ *for* $t \in [t_1, t_2]$ *in terms of* $\max_{t\in[t_1,t_2]}|\omega|_2(t)$ *and* $|\omega|_{\frac{6.02}{3}}(t_1)$*; then derive the upper bound on* $|\bar\nabla\bar\omega|_{3.01}(t)$ *for* $t \in [t_1, t_2]$ *in terms of* $\max_{t\in[t_1,t_2]}|\omega|_2(t)$*,* $|\omega|_{\frac{6.02}{3}}(t_1)$ *and* $|\bar\nabla\bar\omega|_2(t_1)$*; then we derive the upper bound on* $|\omega|_\infty(t)$ *for* $t \in [t_1, t_2]$ *in terms of* $\max_{t\in[t_1,t_2]}|\omega|_2(t)$*,* $|\omega|_{\frac{6.02}{3}}(t_1)$*,* $|\bar\nabla\bar\omega|_2(t_1)$ *and* $\bar\nabla\bar\omega|_{3.01}(t_1)$*. These estimates are used to improve the estimate on* $|\omega|_\infty(t)$ *and to show that* $|\omega|_\infty(t) \le \max(|\omega|_\infty(t_1), e^e)$*, provided that certain assumptions hold. Repeated use of Local Existence and Uniqueness results together with the upper bounds on* $|\omega|_\infty(t)$ *gives the existence and uniqueness of the solution if* $|\omega|_\infty(0) < \infty$ *or if* $|\omega|_4(0) < \infty$*.*

Introduction and statement of results.

The Navier-Stokes Equations for velocity can be written as

$$\partial_t\bar u(\bar x,t) - \nu\Delta\bar u(\bar x,t) + (\bar u(\bar x,t)\cdot\bar\nabla)\bar u(\bar x,t) + \bar\nabla p(\bar x,t) = f(\bar x,t), \quad (1)$$
$$\bar\nabla\cdot\bar u(\bar x,t) = 0,\ \bar x \in \mathbf{R}^3,\ t\in[0,\infty).$$

We assume that $f(\bar x,t) = 0$ and that $\lim_{|\bar x|\to\infty}|\bar u(\bar x,t)| = 0$.

Introducing vorticity $\bar\omega(\bar x,t) = \bar\nabla\times\bar u(\bar x,t)$ allows us to obtain the Navier-Stokes Equations without the pressure term.

The Navier-Stokes Equations for vorticity can be written as

$$\partial_t\bar\omega(\bar x,t) - \nu\Delta\bar\omega(\bar x,t) = -(\bar u(\bar x,t)\cdot\bar\nabla)\bar\omega(\bar x,t) + (\bar\omega(\bar x,t)\cdot\bar\nabla)\bar u(\bar x,t). \quad (2)$$
$$\bar\nabla\cdot\bar\omega(\bar x,t) = 0,\ \bar x\in\mathbf{R}^3,\ t\in[0,\infty).$$

We assume that $\lim_{|x|\to\infty}|\bar\omega(\bar x,t)| = 0$.

The question of existence and uniqueness of a strong solution of these equations in 3 dimensions is a longstanding problem. The solution of this problem is described in this paper.

The main result of the paper are the following Theorems:

Theorem 1: *Suppose that* $|\omega|_2(t) < \infty$ *for* $t \in [t_1, t_2]$ *and also that* $|\omega|_\infty(t_1) < \infty$ *and* $|\bar{\nabla}\bar{\omega}|_{3.01}(t_1) < \infty$. *Then* $|\omega|_\infty(t) < \infty$ *for* $t \in [t_1, t_2]$ *and an upper bound (equation (43)) on* $|\omega|_\infty(t)$ *can be given as a function of* t, $\max_{t\in[t_1,t_2]}|\omega|_2(t)$, $|\omega|_\infty(t_1)$ *and* $|\bar{\nabla}\bar{\omega}|_{3.01}(t_1)$.

Theorem 2: *Suppose that* $|\omega|_2(t) < \infty$ *for* $t \in [t_1, t_2]$ *and that* $|\omega|_\infty(t_1) < \infty$ *and* $|\bar{\nabla}\bar{\omega}|_{3.01}(t_1) < \infty$. *Suppose also that there exists* n_0 *such that for all* $n > n_0$, *either* $|\omega|_n(t) \geq e^e$ *for all* $t \in [t_1, t_2]$ *or* $|\omega|_n(t)$ *is a continuous function of* t *for all* $t \in [t_1, t_2]$. *Then for all sufficiently large* n, $|\omega|_n(t) \leq \max(e^{\frac{1}{n}e^{e^{t_2-t_1}}}|\omega|_n(t_1), e^e)$ *and* $|\omega|_\infty(t) \leq \max(|\omega|_\infty(t_1), e^e)$ *for* $t \in [t_1, t_2]$.

Theorem 3: *Suppose that* $|\omega|_\infty(0) < \infty$. *Then for any* $T > 0$, *there exists a unique solution of the 3D Navier Stokes Equations* $\bar{\omega}(\bar{x}, t)$ *for* $x \in \mathbf{R}^3$ *and* $t \in [0, T]$, *such that* $|\omega|_\infty(t) < \infty$ *for* $t \in [0, T]$.

Theorem 4: *Suppose that* $|\omega|_4(0) < \infty$. *Then for any* $T > 0$, *there exists a unique solution of the 3D Navier Stokes Equations* $\bar{\omega}(\bar{x}, t)$ *for* $x \in \mathbf{R}^3$ *and* $t \in [0, T]$, *such that* $|\omega|_\infty(t) < \infty$ *for* $t \in (0, T]$.

Proof of Theorem 1.

Part 1: Take Navier Stokes equation for $\bar{\omega}$ and take the inner product with $\bar{\omega}(x)$ integrate over space and by parts on the viscosity term to obtain

$$\frac{1}{2}\partial_t \int |\bar{\omega}(\bar{x})|^2 d^3x + \nu \int |\bar{\nabla}\bar{\omega}(\bar{x})|^2 d^3x = \frac{3}{4\pi}\int d^3x \int d^3y (\bar{\omega}(\bar{x}) \cdot \hat{y})(\bar{\omega}(\bar{x}) \times \bar{\omega}(\bar{x}+\bar{y}) \cdot \hat{y})\frac{1}{|y|^3}. \tag{3}$$

The absolute value of the integral on the right hand side of (3) can be estimated as

$$\begin{aligned}
&\left|\int d^3x \int d^3y (\bar{\omega}(\bar{x}) \cdot \hat{y})(\bar{\omega}(\bar{x}) \times \bar{\omega}(\bar{x}+\bar{y}) \cdot \hat{y})\frac{1}{|y|^3}\right| \leq \qquad (4)\\
&\leq \left|\int d^3x \int_{|y|<1} (\bar{\omega}(\bar{x}) \cdot \hat{y})(\bar{\omega}(\bar{x}) \times (\bar{\omega}(\bar{x}+\bar{y}) - \bar{\omega}(\bar{x})) \cdot \hat{y})\frac{d^3y}{|y|^3}\right| + \\
&\qquad + \left|\int d^3x \int_{|y|\geq 1} (\bar{\omega}(\bar{x}) \cdot \hat{y})(\bar{\omega}(\bar{x}) \times \bar{\omega}(\bar{x}+\bar{y}) \cdot \hat{y})\frac{d^3y}{|y|^3}\right| \\
&\leq 3\sqrt{6}\int_0^1 ds \int d^3x \int_{|y|<1} |\bar{\omega}(\bar{x})|^2 |\bar{\nabla}\bar{\omega}(\bar{x}+s\bar{y})|\frac{d^3y}{|y|^2} \\
&\qquad +3\sqrt{2}\int d^3x |\bar{\omega}(\bar{x})|^2 \int_{|y|\geq 1} \frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^3} d^3y
\end{aligned}$$

$$\leq 3\sqrt{6}\int_0^1 ds\int d^3x'|\bar{\nabla}\bar{\omega}(\bar{x}')|\int_{|y|<1} d^3y\frac{|\bar{\omega}(\bar{x}'-s\bar{y})|^2}{|y|^2}+$$

$$+3\sqrt{2}\int d^3x|\bar{\omega}(\bar{x})|^2\int_{|y|\geq 1} d^3y\frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^3}$$

$$\leq 3\sqrt{6}\int_0^1 ds\sqrt{\int d^3x'|\bar{\nabla}\bar{\omega}(\bar{x}')|^2}\times$$

$$\sqrt{\int d^3x'\int_{|y|<1}\frac{|\bar{\omega}(\bar{x}'-s\bar{y})|^2}{|y|^2}d^3y\int_{|z|<1}\frac{|\bar{\omega}(\bar{x}'-s\bar{z})|^2}{|z|^2}d^3z}+\sqrt{24\pi}\int d^3x|\bar{\omega}(\bar{x})|^2(\int d^3y|\bar{\omega}(\bar{y})|^2)^{\frac{1}{2}}$$

$$\leq 3\sqrt{6}\int_0^1 ds\sqrt{\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2}\sqrt{\int d^3x'\int_{|y|\leq 1}\frac{|\bar{\omega}(\bar{x}'-s\bar{y})|^2}{|y|^{3-\epsilon}}d^3y\int_{|z|\leq 1}\frac{|\bar{\omega}(\bar{x}'-s\bar{z})|^2}{|z|^{1+\epsilon}}d^3z}+$$

$$+\sqrt{24\pi}|\omega|_2^2|\omega|_2$$

$$\leq 3\sqrt{6}\int_0^1 ds\sqrt{\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2}\sqrt{\int d^3x|\bar{\omega}(\bar{x})|^2\int_{|y|\leq 1}\frac{d^3y}{|y|^{3-\epsilon}}\int_{|z|\leq 1}\frac{|\bar{\omega}(\bar{x}+s\bar{y}-s\bar{z})|^2}{|z|^{1+\epsilon}}d^3z}+$$

$$+\sqrt{24\pi}|\omega|_2^2|\omega|_2$$

$$\leq 3\sqrt{6}\int_0^1\frac{ds}{s^{1-\frac{\epsilon}{2}}}\sqrt{\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2}\sqrt{\int d^3x|\bar{\omega}(\bar{x})|^2\int_{|y|\leq 1}\frac{d^3y}{|y|^{3-\epsilon}}\int_{|z'|\leq 1}\frac{|\bar{\omega}(\bar{x}+s\bar{y}-\bar{z}')|^2}{|z'|^{1+\epsilon}}d^3z'}+$$

$$+\sqrt{24\pi}|\omega|_2^2|\omega|_2$$

$$\leq 3\sqrt{6}\int_0^1\frac{ds}{s^{1-\frac{\epsilon}{2}}}\sqrt{\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2}$$

$$\sqrt{\int d^3x|\bar{\omega}(\bar{x})|^2\int_{|y|\leq 1}\frac{d^3y}{|y|^{3-\epsilon}}(\int d^3z'|\bar{\omega}(\bar{x}-\bar{z}'+s\bar{y})|^{\frac{2(3-3\epsilon)}{2-3\epsilon}})^{\frac{2-3\epsilon}{3-3\epsilon}}(\int_{|z'|<1}\frac{d^3z'}{|z'|^{3(1-\epsilon^2)}})^{\frac{1}{3(1-\epsilon)}}}+$$

$$+\sqrt{24\pi}|\omega|_2^2|\omega|_2$$

$$\leq 3\sqrt{6}(\frac{4\pi}{\epsilon})^{\frac{1}{2}}(\frac{4\pi}{3\epsilon^2})^{\frac{1}{6(1-\epsilon)}}\int_0^1\frac{ds}{s^{1-\frac{\epsilon}{2}}}\sqrt{\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2}\sqrt{\int d^3x|\bar{\omega}(\bar{x})|^2(\int d^3z|\bar{\omega}(\bar{z})|^{\frac{2(3-3\epsilon)}{2-3\epsilon}})^{\frac{2-3\epsilon}{3-3\epsilon}}}$$

$$+\sqrt{24\pi}|\omega|_2^2|\omega|_2$$

$$\leq\frac{6\sqrt{6}}{\epsilon}(\frac{4\pi}{\epsilon})^{\frac{1}{2}}(\frac{4\pi}{3\epsilon^2})^{\frac{1}{6(1-\epsilon)}}(\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x)^{\frac{3-2\epsilon}{4(1-\epsilon)}}(\int|\bar{\omega}(\bar{x})|^2d^3x)^{\frac{3-4\epsilon}{4-4\epsilon}}+\sqrt{24\pi}|\omega|_2^2|\omega|_2$$

$$\leq\frac{1-2\epsilon}{4-4\epsilon}(\frac{6\sqrt{6}}{\epsilon})^{\frac{4-4\epsilon}{1-2\epsilon}}(\frac{4\pi}{\epsilon})^{\frac{2-2\epsilon}{1-2\epsilon}}(\frac{4\pi}{3\epsilon^2})^{\frac{2}{3(1-2\epsilon)}}(\frac{2^{0.33}}{\nu})^{\frac{3-2\epsilon}{1-2\epsilon}}(\int d^3x|\bar{\omega}(\bar{x})|^2)^{\frac{3-4\epsilon}{1-2\epsilon}}+$$

$$+\frac{\nu}{2^{0.33}}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2+\sqrt{24\pi}|\omega|_2^2|\omega|_2.$$

Notice that in the above estimate ϵ is arbitrarily small. Using this estimate and introducing ε such that

$$\varepsilon = \frac{2\epsilon}{1-2\epsilon}$$

we obtain the following inequality from (3)

$$\frac{1}{2}\partial_t \int |\bar{\omega}(\bar{x})|^2 d^3x + \nu(1-\frac{1}{2^{0.33}})\int |\bar{\nabla}\bar{\omega}(\bar{x})|^2 d^3x \leq$$
$$\leq \frac{3}{4\pi}\left(\frac{0.331\cdot 6^6\cdot (4\pi)^{\frac{8}{3}}}{\varepsilon^{\frac{22}{3}+\frac{13}{3}\varepsilon}\nu^{3+2\varepsilon}}(\int |\bar{\omega}(\bar{x})|^2 d^3x)^{3+\varepsilon} + \sqrt{24\pi}|\omega|_2^3\right). \tag{5}$$

Note that ε can be chosen arbitrarily small. Below we assume that $\varepsilon = \frac{1}{100}$.

Since $\int |\bar{\omega}(\bar{x})|^2 d^3x < \infty$ for $t \in [t_1, t_2]$, there exists $N > 0$ such that $\int |\bar{\omega}(\bar{x})|^2 d^3x(t) \leq N$ for $t \in [t_1, t_2]$. Denote

$$C(t_1,t_2) = \int_{t_1}^{t_2} dt \int |\bar{\omega}(\bar{x})|^2 d^3x, C(t) = \int_0^t ds \int |\bar{\omega}(\bar{x})|^2 d^3x(s).$$

Then we obtain

$$\frac{1}{2}\int |\bar{\omega}(\bar{x})|^2 d^3x(t_2) - \frac{1}{2}\int |\bar{\omega}(\bar{x})|^2 d^3x(t_1) + \nu(1-\frac{1}{2^{0.33}})\int_{t_1}^{t_2} dt \int |\bar{\nabla}\bar{\omega}(\bar{x})|^2 d^3x$$
$$\leq \frac{2.74\cdot 10^6}{\varepsilon^{\frac{22}{3}+\frac{13}{3}\varepsilon}\nu^{3+2\varepsilon}} N^{2+\varepsilon}C(t_1,t_2) + 2.1N^{\frac{1}{2}}C(t_1,t_2) \leq \frac{2.35\cdot 10^{21}}{\nu^{3.02}}N^{2+\varepsilon}C(t_1,t_2) + 2.1N^{\frac{1}{2}}C(t_1,t_2).$$

Therefore

$$\int_{t_1}^{t_2} dt \int d^3x |\bar{\nabla}\bar{\omega}(\bar{x})|^2 \leq \frac{1.075\cdot 10^{22}}{\nu^{4+2\varepsilon}}N^{2+\varepsilon}C(t_1,t_2) + \frac{9.6}{\nu}N^{\frac{1}{2}}C(t_1,t_2) + \frac{4.57}{\nu}N. \tag{6}$$

Part 2: Take the inner product of the Navier-Stokes equation for $\bar{\omega}(\bar{x})$ with $|\omega(\bar{x})|^{\frac{2}{3}\varepsilon}\bar{\omega}(\bar{x})$, integrate over space and integrate the viscosity term by parts to obtain

$$\frac{1}{2+\frac{2}{3}\varepsilon}\partial_t \int |\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x})d^3x + \nu\int \bar{\nabla}(|\omega|^{\frac{2}{3}\varepsilon}(\bar{x})\bar{\omega}(\bar{x}))(\bar{\nabla}\bar{\omega}(\bar{x}))d^3x \tag{7}$$
$$= \frac{3}{4\pi}\int\int (|\omega|^{\frac{2}{3}\varepsilon}(\bar{x})\bar{\omega}(\bar{x})\cdot\hat{y})(\bar{\omega}(\bar{x})\times\bar{\omega}(\bar{x}+\bar{y})\cdot\hat{y})\frac{d^3xd^3y}{|y|^3}.$$

The absolute value of the integral on the right hand side of (7) can be estimated from above as

$$\left|\int\int (|\omega|^{\frac{2}{3}\varepsilon}(\bar{x})\bar{\omega}(\bar{x})\cdot\hat{y})(\bar{\omega}(\bar{x})\times\bar{\omega}(\bar{x}+\bar{y})\cdot\hat{y})\frac{d^3xd^3y}{|y|^3}\right| \leq \tag{8}$$

$$\leq \left| \int\int_{|y|<1} (|\omega|^{\frac{2}{3}\varepsilon}(\bar{x})\bar{\omega}(\bar{x})\cdot\hat{y})(\bar{\omega}(\bar{x}) \times (\bar{\omega}(\bar{x}+\bar{y}) - \bar{\omega}(\bar{x}))\cdot\hat{y}) \frac{d^3x d^3y}{|y|^3} \right| +$$

$$+ \left| \int\int_{|y|\geq 1} (|\omega|^{\frac{2}{3}\varepsilon}(\bar{x})\bar{\omega}(\bar{x})\cdot\hat{y})(\bar{\omega}(\bar{x}) \times \bar{\omega}(\bar{x}+\bar{y})\cdot\hat{y}) \frac{d^3x d^3y}{|y|^3} \right| \leq$$

$$\leq 3\sqrt{6}\int_0^1 ds \int d^3x |\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}) \int_{|y|<1} \frac{|\bar{\nabla}\bar{\omega}(\bar{x}+s\bar{y})|}{|y|^2} d^3y +$$

$$+3\sqrt{2}\int\int_{|y|\geq 1} |\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x})|\omega|(\bar{x}+\bar{y}) \frac{d^3x d^3y}{|y|^3} \leq$$

$$\leq 3\sqrt{6}\int_0^1 ds \int d^3x' |\bar{\nabla}\bar{\omega}(\bar{x}')| \int_{|y|<1} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}'-s\bar{y})}{|y|^2} d^3y +$$

$$+3\sqrt{2}\int |\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}) d^3x \int_{|y|\geq 1} \frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^3} d^3y \leq$$

$$\leq 3\sqrt{6}\int_0^1 ds \sqrt{\int d^3x' \int_{|y|<1} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}'-s\bar{y})}{|y|^2} d^3y \int_{|z|<1} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}'-s\bar{z})}{|z|^2} d^3z} |\bar{\nabla}\bar{\omega}|_2$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2$$

$$\leq 3\sqrt{6}\int_0^1 ds \sqrt{\int d^3x' \int_{|y|\leq 1} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}'-s\bar{y})}{|y|^{3-\varepsilon}} d^3y \int_{|z|\leq 1} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}'-s\bar{z})}{|z|^{1+\varepsilon}} d^3z} \sqrt{\int |\bar{\nabla}\bar{\omega}(\bar{x})|^2 d^3x}$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2$$

$$\leq 3\sqrt{6}\int_0^1 ds \sqrt{\int |\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}'') d^3x'' \int_{|y|\leq 1} \frac{d^3y}{|y|^{3-\varepsilon}} \int_{|z|\leq 1} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}''-s(\bar{z}-\bar{y}))}{|z|^{1+\varepsilon}} d^3z}\ |\bar{\nabla}\bar{\omega}|_2$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2 \leq$$

$$\leq 3\sqrt{6}\int_0^1 ds \sqrt{\int d^3x'' |\omega|^{2+\frac{2}{3}\varepsilon}(x'') \int_{|y|\leq 1} \frac{d^3y}{|y|^{3-\varepsilon}} \int_{|z'|\leq s} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(x''+sy-z')}{|z'|^{1+\varepsilon}} \frac{d^3z'}{s^{2-\varepsilon}}} |\bar{\nabla}\bar{\omega}|_2$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2 \leq$$

$$\leq 3\sqrt{6}\int_0^1 \frac{ds}{s^{1-\frac{\varepsilon}{2}}} \sqrt{\int d^3x |\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x}) \int_{|y|\leq 1} \frac{d^3y}{|y|^{3-\varepsilon}} \int_{|z'|\leq 1} \frac{|\omega|^{2+\frac{2}{3}\varepsilon}(x+sy-z')}{|z'|^{1+\varepsilon}} d^3z'} |\bar{\nabla}\bar{\omega}|_2 +$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2 \leq$$

$$\leq 3\sqrt{6}(\frac{4\pi}{3\varepsilon^2})^{\frac{1}{6(1-\varepsilon)}}$$

$$\int_0^1 \frac{ds}{s^{1-\frac{\varepsilon}{2}}}\sqrt{\int d^3x|\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x})\int_{|y|\leq 1}\frac{d^3y}{|y|^{3-\varepsilon}}(\int |\omega|^{(2+\frac{2}{3}\varepsilon)\frac{3-3\varepsilon}{2-3\varepsilon}}(x+sy-z')d^3z')^{\frac{2-3\varepsilon}{3-3\varepsilon}}}|\bar{\nabla}\bar{\omega}|_2$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2$$

$$\leq 3\sqrt{6}(\frac{4\pi}{\varepsilon})^{\frac{1}{2}}(\frac{4\pi}{3\varepsilon^2})^{\frac{1}{6(1-\varepsilon)}}|\bar{\nabla}\bar{\omega}|_2$$

$$\int_0^1 \frac{ds}{s^{1-\frac{\varepsilon}{2}}}\sqrt{(\int |\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x)^{1+\frac{(1-\frac{7}{3}\varepsilon+\frac{\varepsilon^2}{3})(1-\frac{3}{2}\varepsilon)}{2(1-\frac{5}{3}\varepsilon+\frac{\varepsilon^2}{4})(1-\varepsilon)}}(\int |\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x)^{\frac{(1+\frac{\varepsilon}{3})(1-\frac{3}{2}\varepsilon)}{2(1-\frac{5}{3}\varepsilon+\frac{\varepsilon^2}{4})(1-\varepsilon)}}} +$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2$$

$$\leq \frac{6\sqrt{6}}{\varepsilon}(\frac{4\pi}{\varepsilon})^{\frac{1}{2}}(\frac{4\pi}{3\varepsilon^2})^{\frac{1}{6(1-\varepsilon)}}(\int |\omega|^{2+\frac{2}{3}\varepsilon}(\bar{x})d^3x)^{\frac{1}{2}+\frac{(1-\frac{7}{3}\varepsilon+\frac{\varepsilon^2}{3})(1-\frac{3}{2}\varepsilon)}{4(1-\frac{5}{3}\varepsilon+\frac{\varepsilon^2}{4})(1-\varepsilon)}}(\int |\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x)^{\frac{1}{2}+\frac{(1+\frac{\varepsilon}{3})(1-\frac{3}{2}\varepsilon)}{4(1-\frac{5}{3}\varepsilon+\frac{\varepsilon^2}{4})(1-\varepsilon)}}$$

$$+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2$$

$$\leq 3.0618\cdot 10^5\left(\int |\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x+1\right)\left(\int |\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x\right)^{\alpha}+\sqrt{24\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}|\omega|_2,$$

where $\alpha = \frac{1}{2}+\frac{1}{4}\frac{(1-\frac{7}{3}\varepsilon+\frac{\varepsilon^2}{3})(1-\frac{3}{2}\varepsilon)}{(1-\frac{5}{3}\varepsilon+\frac{\varepsilon^2}{4})(1-\varepsilon)} \approx 0.747$. Putting the estimates together we obtain

$$\frac{1}{2+\frac{2}{3}\varepsilon}\partial_t\int |\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x+\nu\int \bar{\nabla}(|\bar{\omega}(\bar{x})|^{\frac{2}{3}\varepsilon}\bar{\omega}(\bar{x}))(\bar{\nabla}\bar{\omega}(\bar{x}))d^3x \leq \tag{9}$$

$$\leq \frac{3}{4\pi}3.0618\cdot 10^5\left(\int |\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x\right)^{\alpha}\left(\int |\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x+1\right)+\frac{3\sqrt{24\pi}}{4\pi}|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}N^{\frac{1}{2}}$$

$$\leq 0.7305\cdot 10^5\left(\int |\omega|^{2+\frac{2}{3}\varepsilon}d^3x\right)^{\alpha}\left(\int |\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x+1\right)+2.1|\omega|_{2+\frac{2}{3}\varepsilon}^{2+\frac{2}{3}\varepsilon}N^{\frac{1}{2}}.$$

Since we can rewrite the viscosity term as shown in the (10) we obtain that the viscosity term is positive definite

$$\int \bar{\nabla}(|\omega|^{\frac{2}{3}\varepsilon}(\bar{x})\bar{\omega}(\bar{x}))(\bar{\nabla}\bar{\omega})d^3x = \frac{\frac{2}{3}\varepsilon}{(1+\frac{\varepsilon}{3})^2}\int (\bar{\nabla}|\omega|^{1+\frac{\varepsilon}{3}}(\bar{x}))^2d^3x+\int |\omega|^{\frac{2}{3}\varepsilon}(\bar{x})(\bar{\nabla}\bar{\omega}(\bar{x}))^2d^3x \tag{10}$$

From (9) we obtain

$$\int |\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x(t_2) \leq \tag{11}$$

$$
\begin{aligned}
&\le e^{4.22\ N^{\frac{1}{2}}(t_2-t_1)}\left(0.371\cdot 10^5\int_{t_1}^{t_2}dt\left(\int|\bar{\nabla}\bar{\omega}|^2(\bar{x})d^3x+1\right)+\left(\int|\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x(t_1)\right)^{1-\alpha}\right)^{\frac{1}{1-\alpha}}\\
&\le e^{4.22N^{\frac{1}{2}}(t_2-t_1)}\\
&\left(\frac{3.99\cdot 10^{26}}{\nu^{4+2\varepsilon}}N^{2+\varepsilon}C(t_1,t_2)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t_1,t_2)+1.7\cdot 10^5\frac{N}{\nu}+0.371\cdot 10^5(t_2-t_1)+\right.\\
&\qquad\qquad\left.+1+|\omega|_{2+\frac{2}{3}\varepsilon}^{(1-\alpha)(2+\frac{2}{3}\varepsilon)}(t_1)\right)^{\frac{1}{1-\alpha}}
\end{aligned}
$$

Therefore we obtain that

$$
\begin{aligned}
&\nu\int_{t_1}^{t_2}dt\int\bar{\nabla}(|\bar{\omega}(\bar{x})|^{\frac{2}{3}\varepsilon}\bar{\omega}(\bar{x}))(\bar{\nabla}\bar{\omega}(\bar{x}))d^3x\le \qquad (12)\\
&\le 0.7305\cdot 10^5\max_{t\in[t_1,t_2]}\left(\int|\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x(t)\right)^{\alpha}\int_{t_1}^{t_2}dt\left(\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x+1\right)+\\
&\qquad\qquad+\frac{1}{2+\frac{2}{3}\varepsilon}\left(\int|\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x(t_1)+\int|\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3x(t_2)\right)+\\
&\qquad\qquad\qquad+2.1\max_{t\in[t_1,t_2]}\int|\bar{\omega}(\bar{x})|^{2+\frac{2}{3}\varepsilon}d^3xN^{\frac{1}{2}}(t_2-t_1)\\
&\le 1.97e^{4.22N^{\frac{1}{2}}(t_2-t_1)}\cdot\left(1+\frac{3.99\cdot 10^{26}}{\nu^{4+2\varepsilon}}N^{2+\varepsilon}C(t_1,t_2)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t_1,t_2)+\right.\\
&\qquad\qquad\left.+1.7\cdot 10^5\frac{N}{\nu}+0.371\cdot 10^5(t_2-t_1)+1.066N^{\frac{1}{2}}(t_2-t_1)+|\omega|_{2+\frac{2}{3}\varepsilon}^{(1-\alpha)(2+\frac{2}{3}\varepsilon)}(t_1)\right)^5
\end{aligned}
$$

Therefore

$$
\begin{aligned}
&\frac{\frac{2}{3}\varepsilon}{(1+\frac{\varepsilon}{3})^2}\int_{t_1}^{t_2}dt\int(\bar{\nabla}|\bar{\omega}(\bar{x})|^{1+\frac{\varepsilon}{3}})^2d^3x+\int_{t_1}^{t_2}dt\int|\bar{\omega}(\bar{x})|^{\frac{2}{3}\varepsilon}(\bar{\nabla}\bar{\omega}(\bar{x}))^2d^3x\le \qquad (13)\\
&\le\frac{1.97}{\nu}e^{4.22N^{\frac{1}{2}}(t_2-t_1)}\cdot\left(1+\frac{3.99\cdot 10^{26}}{\nu^{4+2\varepsilon}}N^{2+\varepsilon}C(t_1,t_2)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t_1,t_2)+\right.\\
&\qquad\left.+1.7\cdot 10^5\frac{N}{\nu}+0.371\cdot 10^5(t_2-t_1)+1.066N^{\frac{1}{2}}(t_2-t_1)+|\omega|_{2+\frac{2}{3}\varepsilon}^{(1-\alpha)(2+\frac{2}{3}\varepsilon)}\right)^5.
\end{aligned}
$$

Part 3: Differentiate with respect to x the Navier-Stokes equation for $\bar{\omega}$.

$$
\partial_t\partial_x\bar{\omega}(\bar{x})-\nu\partial_x\Delta\bar{\omega}(\bar{x})=\partial_x\left((\bar{\omega}(\bar{x})\cdot\bar{\nabla})\bar{u}(\bar{x})\right)-\partial_x\left((\bar{u}(\bar{x})\cdot\bar{\nabla})\bar{\omega}(\bar{x})\right). \qquad (14)
$$

Take inner product with $\partial_x\bar{\omega}$

$$
\frac{1}{2}\partial_t(\partial_x\bar{\omega}(\bar{x}))^2-\nu\partial_x\bar{\omega}(\bar{x})\cdot\partial_x\Delta\bar{\omega}(\bar{x})=\partial_x\bar{\omega}(\bar{x})\cdot\partial_x\left((\bar{\omega}(\bar{x})\cdot\bar{\nabla})\bar{u}(\bar{x})\right)-\partial_x\bar{\omega}(\bar{x})\cdot\partial_x\left((\bar{u}(\bar{x})\cdot\bar{\nabla})\bar{\omega}(\bar{x})\right)
$$

Integrate over space and by parts.

$$\frac{1}{2}\partial_t\int(\partial_x\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{xx}\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{xy}\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{xz}\bar\omega(\bar x))^2d^3x=$$
$$=-\int\partial_{xx}\bar\omega(\bar x)\cdot(\bar\omega(\bar x)\cdot\bar\nabla)\bar u(\bar x)d^3x+\int\partial_{xx}\bar\omega\cdot(\bar u(\bar x)\cdot\bar\nabla)\bar\omega(\bar x)d^3x. \tag{15}$$

Add the equations obtained similarly but with differentiation with respect to y or z instead of x.

$$\frac{1}{2}\partial_t\int(\partial_y\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{yy}\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{xy}\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{yz}\bar\omega(\bar x))^2d^3x=$$
$$=-\int\partial_{yy}\bar\omega(\bar x)\cdot(\bar\omega(\bar x)\cdot\bar\nabla)\bar u(\bar x)d^3x+\int\partial_{yy}\bar\omega(\bar x)\cdot(\bar u(\bar x)\cdot\bar\nabla)\bar\omega(\bar x)d^3x. \tag{16}$$

and

$$\frac{1}{2}\partial_t\int(\partial_z\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{zz}\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{xz}\bar\omega(\bar x))^2d^3x+\nu\int(\partial_{yz}\bar\omega(\bar x))^2d^3x=$$
$$=-\int\partial_{zz}\bar\omega(\bar x)\cdot(\bar\omega(\bar x)\cdot\bar\nabla)\bar u(\bar x)d^3x+\int\partial_{zz}\bar\omega(\bar x)\cdot(\bar u(\bar x)\cdot\bar\nabla)\bar\omega(\bar x)d^3x. \tag{17}$$

to get

$$\frac{1}{2}\partial_t\int|\nabla\omega|^2(\bar x)d^3x+\nu\int|\bar\nabla\bar\nabla\omega|^2(\bar x)d^3x= \tag{18}$$
$$=-\int\Delta\bar\omega(\bar x)\cdot(\bar\omega(\bar x)\cdot\bar\nabla)\bar u(\bar x)d^3x+\int\Delta\bar\omega(\bar x)\cdot(\bar u(\bar x)\cdot\bar\nabla)\bar\omega(\bar x)d^3x.$$

Plugging the expression for $\bar u$ in terms of $\bar\omega$

$$\bar u(\bar x)=\frac{1}{4\pi}\int\frac{\bar\omega(\bar y)\times(\bar x-\bar y)}{|x-y|^3}d^3y=\frac{1}{4\pi}\int\frac{\bar\omega(\bar x+\bar y)\times\hat y}{|y|^2}d^3y$$

on the right hand side, we obtain

$$\frac{1}{2}\partial_t\int|\bar\nabla\bar\omega(\bar x)|^2d^3x+\nu\int|\bar\nabla\bar\nabla\bar\omega(\bar x)|^2d^3x=-\frac{1}{4\pi}\int\int\Delta\bar\omega(\bar x)\cdot(\bar\omega(\bar x)\cdot\bar\nabla)\frac{\bar\omega(\bar x+\bar y)\times\hat y}{|y|^2}d^3xd^3y$$
$$+\frac{1}{4\pi}\int\int\Delta\bar\omega(\bar x)\cdot\frac{((\bar\omega(\bar x+\bar y)\times\hat y)\cdot\bar\nabla)\bar\omega(\bar x)}{|y|^2}d^3xd^3y. \tag{19}$$

We can rewrite the right hand side of (19) as

$$-\frac{1}{4\pi}\int\int_{|y|\leq1}\Delta\bar\omega(\bar x)\cdot(\bar\omega(\bar x)\cdot\bar\nabla)\frac{\bar\omega(\bar x+\bar y)\times\hat y}{|y|^2}d^3xd^3y- \tag{20}$$

$$-\frac{1}{4\pi}\int\int_{|y|>1}\Delta\bar{\omega}(\bar{x})\cdot(\bar{\omega}(\bar{x})\cdot\bar{\nabla})\frac{\bar{\omega}(\bar{x}+\bar{y})\times\hat{y}}{|y|^2}d^3xd^3y+$$
$$+\frac{1}{4\pi}\int\int_{|y|\leq 1}\Delta\bar{\omega}(\bar{x})\cdot\frac{((\bar{\omega}(\bar{x}+\bar{y})\times\hat{y})\cdot\bar{\nabla})\bar{\omega}(\bar{x})}{|y|^2}d^3xd^3y+$$
$$+\frac{1}{4\pi}\int\int_{|y|>1}\Delta\bar{\omega}(\bar{x})\cdot\frac{((\bar{\omega}(\bar{x}+\bar{y})\times\hat{y})\cdot\bar{\nabla})\bar{\omega}(\bar{x})}{|y|^2}d^3xd^3y.$$

The absolute value of the integrals over $|y|\leq 1$ can be estimated from above by

$$\frac{3}{2\pi}\int\int_{|y|\leq 1}|\Delta\bar{\omega}(\bar{x})||\bar{\omega}(\bar{x})|\frac{|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}d^3xd^3y+\frac{3}{2\pi}\int\int_{|y|\leq 1}|\Delta\bar{\omega}(\bar{x})||\bar{\omega}(\bar{x}+\bar{y})|\frac{|\bar{\nabla}\bar{\omega}(\bar{x})|}{|y|^2}d^3xd^3y$$
$$=\frac{3}{2\pi}\int\int_{|y|\leq 1}|\Delta\bar{\omega}(\bar{x}-\bar{y})||\bar{\omega}(\bar{x}-\bar{y})|\frac{|\bar{\nabla}\bar{\omega}(\bar{x})|}{|y|^2}d^3xd^3y$$
$$+\frac{3}{2\pi}\int\int_{|y|\leq 1}|\Delta\bar{\omega}(\bar{x})||\bar{\omega}(\bar{x}+\bar{y})|\frac{|\bar{\nabla}\bar{\omega}(\bar{x})|}{|y|^2}d^3xd^3y$$
$$=\frac{3}{2\pi}\int_{|y|\leq 1}\frac{d^3y}{|y|^2}\int|\Delta\bar{\omega}(\bar{x}-\bar{y})||\bar{\omega}(\bar{x}-\bar{y})||\bar{\nabla}\bar{\omega}(\bar{x})|d^3x+$$
$$+\frac{3}{2\pi}\int|\Delta\bar{\omega}(\bar{x})||\bar{\nabla}\bar{\omega}(\bar{x})|\int_{|y|\leq 1}\frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}d^3y$$
$$\leq\frac{3}{4\pi}\int_{|y|\leq 1}\frac{d^3y}{|y|^2}\int(\frac{\nu}{5}|\Delta\bar{\omega}(\bar{x}-\bar{y})|^2+\frac{5}{\nu}|\bar{\omega}(\bar{x}-\bar{y})|^2|\bar{\nabla}\bar{\omega}(\bar{x})|^2)d^3x+$$
$$+\frac{3}{4\pi}\int(\frac{\nu}{5}|\Delta\bar{\omega}(\bar{x})|^2+\frac{5}{\nu}|\bar{\nabla}\bar{\omega}(\bar{x})|^2\int_{|y|\leq 1}\frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}d^3y\int_{|z|\leq 1}\frac{|\bar{\omega}(\bar{x}+\bar{z})|}{|z|^2}d^3z)d^3x$$
$$=\frac{3}{4\pi}\frac{\nu}{5}\int|\Delta\bar{\omega}(\bar{x})|^2(1+\int_{|y|\leq 1}\frac{d^3y}{|y|^2})+\frac{3}{4\pi}\frac{5}{\nu}\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x\int_{|y|\leq 1}\frac{|\bar{\omega}(\bar{x}+\bar{y})|^2}{|y|^2}d^3y(\int\frac{d^3z}{|z|^2}+1)$$
$$\leq\frac{3}{4\pi}\frac{\nu}{5}\int|\Delta\bar{\omega}(\bar{x})|^2d^3x(1+\int_{|y|\leq 1}\frac{d^3y}{|y|^2})+$$
$$+\frac{3}{4\pi}\frac{5}{\nu}\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x(\int|\omega|^{6+2\varepsilon}(\bar{x})d^3x)^{\frac{1}{3+\varepsilon}}(4\pi+1)(4\pi)^{\frac{2+\varepsilon}{3+\varepsilon}}(\frac{2+\varepsilon}{\varepsilon})^{\frac{2+\varepsilon}{3+\varepsilon}}$$
$$\leq\frac{3(1+4\pi)}{4\pi}\frac{\nu}{5}\int|\Delta\omega(\bar{x})|^2d^3x+\frac{30}{\nu\varepsilon}(4\pi+1)\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x(\int|\bar{\nabla}|\bar{\omega}|^{1+\frac{\varepsilon}{3}}(\bar{x})|^2d^3x)^{\frac{3}{3+\varepsilon}}.$$

The absolute value of the integral over $|y|>1$ is bounded by

$$\frac{1}{4\pi}\int\int_{|y|>1}\left|\Delta\bar{\omega}(\bar{x})\cdot(\bar{\omega}(\bar{x})\cdot\bar{\nabla})\frac{\bar{\omega}(\bar{x}+\bar{y})\times\hat{y}}{|y|^2}\right|d^3xd^3y+ \tag{21}$$
$$+\frac{1}{4\pi}\int\int_{|y|>1}\left|\Delta\bar{\omega}(\bar{x})\cdot\frac{((\bar{\omega}(\bar{x}+\bar{y})\times\hat{y})\cdot\bar{\nabla})\bar{\omega}(\bar{x})}{|y|^2}\right|d^3xd^3y$$

$$\begin{aligned}&\leq \frac{3}{2\pi}\int\int_{|y|>1}|\Delta\bar{\omega}(\bar{x})||\bar{\omega}(\bar{x})||\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|d^3x\frac{d^3y}{|y|^2}+\\ &\qquad\qquad\qquad +\frac{3}{2\pi}\int|\Delta\bar{\omega}(\bar{x})||\bar{\nabla}\bar{\omega}(\bar{x})|\int_{|y|\geq 1}\frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}d^3yd^3x\\ &\leq \frac{3}{4\pi}\frac{\nu}{5}\int|\Delta\bar{\omega}(\bar{x})|^2d^3x+\frac{15}{\nu}\int|\omega|^2(\bar{x})d^3x(\int|\bar{\nabla}\bar{\omega}|^2(\bar{x}+\bar{y})d^3y)+\\ &\qquad\qquad +\frac{3}{4\pi}\frac{\nu}{5}\int|\Delta\bar{\omega}(\bar{x})|^2d^3x+\frac{15}{\nu}\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2(\int|\bar{\omega}(\bar{x}+\bar{y})|^2d^3y)d^3x.\end{aligned}$$

Therefore we obtain from (19) that

$$\begin{aligned}&\frac{1}{2}\partial_t\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x+\frac{(8\pi-9)\nu}{20\pi}\int|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x\leq \qquad (22)\\ &\leq\frac{30(4\pi+1)}{\varepsilon\nu}\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x(\int(\bar{\nabla}|\bar{\omega}(\bar{x})|^{1+\frac{\varepsilon}{3}})^2d^3x)^{\frac{3}{3+\varepsilon}}+\frac{30}{\nu}\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x\int|\bar{\omega}(\bar{x})|^2d^3x.\end{aligned}$$

We derive that

$$\partial_t\ln\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x\leq\frac{816}{\varepsilon\nu}(\int|\bar{\nabla}|\bar{\omega}(\bar{x})|^{1+\frac{\varepsilon}{3}}|^2d^3x+1)+\frac{60}{\nu}\int|\bar{\omega}(\bar{x})|^2d^3x. \qquad (23)$$

Therefore

$$\begin{aligned}&\ln\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x(t_2)-\ln\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x(t_1)\leq\\ &\leq\frac{8.16\cdot 10^4}{\nu}\left(\int_{t_1}^{t_2}dt\int|\bar{\nabla}|\bar{\omega}(\bar{x})|^{1+\frac{\varepsilon}{3}}|^2d^3x(t)+(t_2-t_1)\right)+\frac{60}{\nu}C(t_1,t_2)\\ &\leq\frac{8.16\cdot 10^4}{\nu}(t_2-t_1)+\frac{60}{\nu}C(t_1,t_2)+\frac{2.432\cdot 10^7}{\nu^2}e^{4.22N^{\frac{1}{2}}(t_2-t_1)}\times \qquad (24)\\ &\left(1+\frac{3.99\cdot 10^{26}}{\nu^{4.02}}N^{2.01}C(t_1,t_2)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t_1,t_2)+1.7\cdot 10^5\frac{N}{\nu}+0.371\cdot 10^5(t_2-t_1)\right.\\ &\qquad\qquad\qquad\left.+1.066N^{\frac{1}{2}}(t_2-t_1)+|\bar{\omega}|_{2+\frac{2}{3}\varepsilon}^{(2+\frac{2}{3}\varepsilon)(1-\alpha)}\right)^5\end{aligned}$$

This implies

$$\begin{aligned}&\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x(t_2)\leq\int|\bar{\nabla}\bar{\omega}(\bar{x})|^2d^3x(t_1)\exp\left(\frac{8.16\cdot 10^4}{\nu}(t_2-t_1)+\frac{60}{\nu}C(t_1,t_2)+\right. \qquad (25)\\ &+\frac{2.432\cdot 10^7}{\nu^2}e^{4.22N^{\frac{1}{2}}(t_2-t_1)}\left(1+\frac{3.99\cdot 10^{26}}{\nu^{4.02}}N^{2.01}C(t_1,t_2)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t_1,t_2)+\right.\\ &\left.\left.+1.7\cdot 10^5\frac{N}{\nu}+0.371\cdot 10^5(t_2-t_1)+1.066N^{\frac{1}{2}}(t_2-t_1)+|\omega|_{2+\frac{2}{3}\varepsilon}^{(2+\frac{2}{3}\varepsilon)(1-\alpha)}(t_1)\right)^5\right)\end{aligned}$$

We obtain

$$\begin{aligned}\int_{t_1}^{t_2} dt \int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x \le \frac{8.16\ \pi\ 10^5}{(8\pi-9)\nu^2}\int_{t_1}^{t_2} dt(\int |\bar\nabla\bar\omega(\bar x)|^2 d^3x)(\int (\bar\nabla|\bar\omega(\bar x)|^{1+\frac{\varepsilon}{3}})^2 d^3x)^{\frac{3}{3+\varepsilon}} \\ +\frac{10\pi}{(8\pi-9)\nu}\int |\bar\nabla\bar\omega(\bar x)|^2 d^3x(t_1) + \frac{10\pi}{(8\pi-9)\nu}\int |\bar\nabla\bar\omega(\bar x)|^2 d^3x(t_1) + \\ +\frac{6\cdot 10^2\pi}{(8\pi-9)\nu}\int_{t_1}^{t_2} dt\int |\bar\nabla\bar\omega(\bar x)|^2 d^3x\int |\bar\omega(\bar x)|^2 d^3x \\ \le \frac{1.945}{\nu}\int |\bar\nabla\bar\omega(\bar x)|^2 d^3x(t_1)\ \exp\Bigg(\frac{1.632\cdot 10^5(t_2-t_1)}{\nu} + 120\frac{C(t_1,t_2)}{\nu} + \\ +\frac{4.864\cdot 10^7}{\nu^2}e^{4.22N^{\frac{1}{2}}(t_2-t_1)}\cdot\Bigg(1+\frac{3.99\cdot 10^{26}}{\nu^{4.02}}N^{2+\varepsilon}C(t_1,t_2)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t_1,t_2)+ \\ +1.7\cdot 10^5\frac{N}{\nu}+0.371\cdot 10^5(t_2-t_1)+1.066N^{\frac{1}{2}}(t_2-t_1)+|\omega|_{2+\frac{2}{3}\varepsilon}^{(2+\frac{2}{3}\varepsilon)(1-\alpha)}(t_1)\Bigg)^5\Bigg).\end{aligned}$$

Part 4: Now we are going to estimate $\int |\bar\nabla\bar\omega|^{3+\varepsilon}(\bar x)d^3x$.

Taking the gradient of the Navier-Stokes equation for ω and then taking the scalar product with $\bar\nabla\bar\omega$ we obtain

$$\begin{aligned}\frac{1}{2}\partial_t(\bar\nabla\bar\omega)^2 - \nu\sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i(\Delta\bar\omega(\bar x)) = \sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i\left((\bar\omega(\bar x)\cdot\bar\nabla)\bar u(\bar x)\right) - \\ -\sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i\left((\bar u(\bar x)\cdot\bar\nabla)\bar\omega(\bar x)\right).\end{aligned} \tag{26}$$

Multiplying (26) by $|\bar\nabla\bar\omega|^{1+\varepsilon}$ and integrating we obtain

$$\begin{aligned}\frac{1}{3+\varepsilon}\partial_t\int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x - \nu\int |\bar\nabla\bar\omega|^{1+\varepsilon}(\bar x)\sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i(\Delta\bar\omega(\bar x))d^3x \\ = \int |\bar\nabla\bar\omega(\bar x)|^{1+\varepsilon}\sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i\left((\bar\omega(\bar x)\cdot\bar\nabla)\int\frac{\bar\omega(\bar x+\bar y)\times\hat y}{|y|^2}\right)d^3y d^3x - \\ -\int |\bar\nabla\bar\omega(\bar x)|^{1+\varepsilon}\sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i\left((\int\frac{\bar\omega(\bar x+\bar y)\times\hat y}{|y|^2}d^3y\cdot\bar\nabla)\bar\omega(\bar x)\right)d^3x.\end{aligned} \tag{27}$$

The viscosity term in (27) can be rewritten as

$$\begin{aligned}-\nu\int d^3x|\bar\nabla\bar\omega(\bar x)|^{1+\varepsilon}\sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i(\sum_j \partial_j\partial_j\bar\omega(\bar x)) = \\ = \nu\int d^3x|\bar\nabla\bar\omega(\bar x)|^{1+\varepsilon}\sum_i\sum_j(\partial_i\partial_j\bar\omega(\bar x))^2 + \nu\int d^3x\frac{1+\varepsilon}{|\bar\nabla\bar\omega(\bar x)|^{1-\varepsilon}}\sum_j(\sum_i \partial_i\bar\omega(\bar x)\cdot\partial_i\partial_j\bar\omega(\bar x))^2.\end{aligned} \tag{28}$$

Therefore the viscosity term in the equation (27) is positive definite.

The absolute value of the right hand side of (27) can be rewritten as

$$\begin{aligned}
&\left|\sum_{i,j,k}\int\int d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}\partial_i\omega_k(\bar{x})\partial_i\omega_j(\bar{x})\ \frac{\partial_j(\bar{\omega}(\bar{x}+\bar{y})\times\hat{y})_k}{|y|^2}+\right. \qquad (29)\\
&\quad+\sum_{i,j,k}\int\int d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}\partial_i\omega_k(\bar{x})\omega_j(\bar{x})\frac{\partial_i\partial_j(\bar{\omega}(\bar{x}+\bar{y})\times\hat{y})_k}{|y|^2}-\\
&\qquad-\sum_{i,j,k}\int\int d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}\partial_i\omega_k(\bar{x})\frac{\partial_i(\bar{\omega}(\bar{x}+\bar{y})\times\hat{y})_j}{|y|^2}\ \partial_j\omega_k(\bar{x})-\\
&\qquad\quad\left.-\sum_{i,j,k}\int\int d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}\partial_i\omega_k(\bar{x})\frac{(\bar{\omega}(\bar{x}+\bar{y})\times\hat{y})_j}{|y|^2}\ \partial_i\partial_j\omega_k(\bar{x})\right|\le\\
&\le 12\sqrt{3}\int\int d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}\frac{|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}+\\
&\qquad+6\sqrt{3}\int d^3x\int_{|y|\le 1}d^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}|\bar{\omega}(\bar{x})|\frac{|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}+\\
&\quad+6\sqrt{3}\int d^3x\int_{|y|>1}d^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}|\bar{\omega}(\bar{x})|\frac{|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}+\\
&\qquad+6\sqrt{3}\int\int_{|y|\le 1}d^3xd^3y|\bar{\nabla}\bar{\omega}|^{2+\varepsilon}(\bar{x})|\bar{\omega}(\bar{x}+\bar{y})|\frac{|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|}{|y|^2}+\\
&\qquad\quad+6\sqrt{3}\int\int_{|y|>1}d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}|\bar{\omega}(\bar{x}+\bar{y})|\frac{|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|}{|y|^2}.
\end{aligned}$$

The first integral on the right hand side of (29) can be estimated as

$$\begin{aligned}
&\int\int_{|y|\le 1}d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}\frac{|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}+\int\int_{|y|>1}d^3xd^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}\frac{|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2} \qquad (30)\\
&\le\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|^6d^3y)^{\frac{1}{6}}(\frac{20\pi}{3})^{\frac{5}{6}}+\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}\int d^3y\frac{|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}\\
&\le\frac{20\pi}{3}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int d^3y|\bar{\nabla}|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})||^2)^{\frac{1}{2}}+\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}\int d^3y\frac{|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}\le\\
&\le\frac{20\pi}{\sqrt{3}}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int d^3y|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{y})|^2)^{\frac{1}{2}}+\sqrt{4\pi}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int d^3y|\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|^2)^{\frac{1}{2}}\le\\
&\le\frac{20\pi}{\sqrt{3}}\int d^3x|\bar{\nabla}\bar{\omega}|^{3+\varepsilon}(\int d^3y|\bar{\nabla}\bar{\nabla}\bar{\omega}|^2+1)+\sqrt{4\pi}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int d^3y|\bar{\nabla}\bar{\omega}(\bar{y})|^2+1).
\end{aligned}$$

The second integral on the right hand side of (29) can be evaluated as

$$\int d^3x \int_{|y|\le 1} d^3y |\bar\nabla\bar\omega(\bar x)|^{2+\varepsilon}|\bar\omega(\bar x)|\frac{|\bar\nabla\bar\nabla\bar\omega(\bar x+\bar y)|}{|y|^2} \tag{31}$$

$$= \int d^3x \int_{|y|\le 1} d^3y |\bar\nabla\bar\omega(\bar x-\bar y)|^{2+\varepsilon}|\bar\omega(\bar x-\bar y)|\frac{|\bar\nabla\bar\nabla\bar\omega(\bar x)|}{|y|^2} \le \sqrt{\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x} \times$$

$$\times\sqrt{\int d^3x \int_{|y|\le 1} d^3y \frac{|\bar\nabla\bar\omega(\bar x-\bar y)|^{2+\varepsilon}|\bar\omega(\bar x-\bar y)|}{|y|^2}\int_{|z|\le 1} d^3z \frac{|\bar\nabla\bar\omega(\bar x-\bar z)|^{2+\varepsilon}|\bar\omega(\bar x-\bar z)|}{|z|^2}}$$

$$\le \sqrt{\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x} \times$$

$$\times\sqrt{\int d^3x \int_{|y|\le 1} d^3y \frac{|\bar\nabla\bar\omega(\bar x-\bar y)|^{2+\varepsilon}|\bar\omega(\bar x-\bar y)|}{|y|^{3-\frac{\varepsilon}{2}}}\int_{|z|\le 1} d^3z \frac{|\bar\nabla\bar\omega(\bar x-\bar z)|^{2+\varepsilon}|\bar\omega(\bar x-\bar z)|}{|z|^{1+\frac{\varepsilon}{2}}}}$$

$$\le \sqrt{\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x}(\int \frac{d^3z}{|z|^{3(1-\frac{\varepsilon^2}{4})}})^{\frac{1}{6(1-\frac{\varepsilon}{2})}} \times$$

$$\sqrt{\int d^3x' |\bar\nabla\bar\omega|^{2+\varepsilon}|\bar\omega(\bar x')| \int_{|y|\le 1}\frac{d^3y}{|y|^{3-\frac{\varepsilon}{2}}}\left(\int d^3z |\bar\nabla\bar\omega(\bar x'+\bar y-\bar z)|^{\frac{3(1-\frac{\varepsilon^2}{4})}{1-\frac{3\varepsilon}{4}}}|\bar\omega(\bar x'+\bar y-\bar z)|^{\frac{3(1-\frac{\varepsilon}{2})}{2(1-\frac{3}{4}\varepsilon)}}\right)^{\frac{2-\frac{3}{2}\varepsilon}{3-\frac{3}{2}\varepsilon}}}$$

$$\le (\frac{16\pi}{3\varepsilon^2})^{\frac{1}{6-3\varepsilon}}(\frac{8\pi}{\varepsilon})^{\frac{1}{2}}\sqrt{\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x}\sqrt{\int d^3x |\bar\omega(\bar x)|^{2+\varepsilon}|\bar\omega(\bar x)|}$$

$$\cdot\left(\int d^3z |\bar\nabla\bar\omega|^{\frac{3(1-\frac{\varepsilon^2}{4})}{1-\frac{3}{4}\varepsilon}}|\bar\omega(\bar z)|^{\frac{3(1-\frac{\varepsilon}{2})}{2(1-\frac{3}{4}\varepsilon)}}\right)^{\frac{1}{3}\frac{1-\frac{3}{4}\varepsilon}{1-\frac{\varepsilon}{2}}}$$

$$\le 629.2(\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x)^{\frac{1}{2}}(\int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x)^{\frac{1}{3}\frac{1+\frac{\varepsilon}{2}}{1+\frac{\varepsilon}{3}}}(\int |\bar\nabla\bar\omega(\bar x)|^2 d^3x)^{\frac{1}{8}\frac{1+\varepsilon}{1+\frac{\varepsilon}{3}}}(\int |\omega|^2(\bar x)d^3x)^{\frac{1}{8}\frac{1-\frac{\varepsilon}{3}}{1+\frac{\varepsilon}{3}}}$$

$$(\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x)^{\frac{1}{4}\frac{1+\frac{3}{2}\varepsilon-\frac{\varepsilon^2}{6}}{1-\frac{7}{6}\varepsilon+\frac{5\varepsilon^2}{18}}\frac{1-\frac{5}{6}\varepsilon}{1-\frac{\varepsilon}{2}}}(\int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x)^{\frac{1}{6}\frac{1-\frac{5}{2}\varepsilon+\frac{\varepsilon^2}{2}}{1-\frac{7}{6}\varepsilon+\frac{5\varepsilon^2}{18}}\frac{1-\frac{5}{6}\varepsilon}{1-\frac{\varepsilon}{2}}}(\int |\bar\nabla\bar\omega|^2(\bar x)d^3x)^{\frac{1}{4}}$$

$$\le 1.1\cdot 10^3(\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x+1)(\int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x+1)(\int |\bar\nabla\bar\omega(\bar x)|^2 d^3x+1)(N+1).$$

The third integral on the right hand side of the equation (27) can be evaluated as

$$\int\int_{|y|>1} |\bar\nabla\bar\omega(\bar x)|^{2+\varepsilon}|\bar\omega(\bar x)|\frac{|\bar\nabla\bar\nabla\bar\omega(\bar x+\bar y)|}{|y|^2}d^3x d^3y \tag{32}$$

$$
\begin{aligned}
&\le \int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}|\bar{\omega}(\bar{x})|\sqrt{\int d^3y|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x}+\bar{y})|^2 d^3y}\sqrt{\int_{|y|>1}\frac{d^3y}{|y|^4}}\\
&\le \sqrt{4\pi}\sqrt{\int d^3y|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{y})|^2}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}|\bar{\omega}(\bar{x})|\\
&\le \sqrt{4\pi}\sqrt{\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2}(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon})^{\frac{2+\varepsilon}{3+\varepsilon}}(\int d^3x|\bar{\omega}(\bar{x})|^{3+\varepsilon})^{\frac{1}{3+\varepsilon}}\\
&\le \sqrt{4\pi}\sqrt{\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2}(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon})^{\frac{2+\varepsilon}{3+\varepsilon}}(\int d^3x|\bar{\omega}(\bar{x})|^6)^{\frac{1+\varepsilon}{4(3+\varepsilon)}}(\int d^3x|\omega|^2(\bar{x}))^{\frac{3-\varepsilon}{4(3+\varepsilon)}}\\
&\le \sqrt{4\pi}\sqrt{\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2}(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon})^{\frac{2+\varepsilon}{3+\varepsilon}}(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2)^{\frac{1+\varepsilon}{4(1+\frac{\varepsilon}{3})}}(\int d^3x|\bar{\omega}(\bar{x})|^2)^{\frac{3-\varepsilon}{4(3+\varepsilon)}}\\
&\le \sqrt{4\pi}\sqrt{\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2}(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon})^{\frac{2+\varepsilon}{3+\varepsilon}}(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2)^{\frac{1+\varepsilon}{4(1+\frac{\varepsilon}{3})}}(\int d^3x|\bar{\omega}(\bar{x})|^2)^{\frac{3-\varepsilon}{4(3+\varepsilon)}}\\
&\le \sqrt{4\pi}(\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}|^2(\bar{x})+1)(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}+1)(\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^2+1)(\int d^3x|\bar{\omega}(\bar{x})|^2+1).
\end{aligned}
$$

The fourth integral in (27) can be estimated as

$$
\begin{aligned}
&\int\int_{|y|\le 1} d^3x d^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}|\bar{\omega}(\bar{x}+\bar{y})|\frac{|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|}{|y|^2} \qquad (33)\\
&= \int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})||\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}(\int_{|y|\le 1} d^3y\frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2})\\
&\le \int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})||\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}(\int d^3y|\bar{\omega}(\bar{x}+\bar{y})|^6)^{\frac{1}{6}}(\int_{|y|\le 1}\frac{d^3y}{|y|^{\frac{12}{5}}})^{\frac{5}{6}}\\
&\le \frac{\nu}{25}\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}+\frac{25}{4\nu}(\frac{20\pi}{3})^{\frac{5}{3}}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int d^3y|\bar{\nabla}\bar{\omega}(\bar{y})|^2)\\
&\le \frac{\nu}{25}\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}+\frac{25}{4\nu}(\frac{20\pi}{3})^{\frac{5}{3}}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}\int d^3y|\bar{\nabla}\bar{\omega}(\bar{y})|^2.
\end{aligned}
$$

The fifth integral in (27) can be estimated as

$$
\begin{aligned}
&\int d^3x\int_{|y|>1} d^3y|\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}|\bar{\omega}(\bar{x}+\bar{y})|\frac{|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|}{|y|^2}\le \qquad (34)\\
&\le \int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{2+\varepsilon}|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|\int d^3y\frac{|\bar{\omega}(\bar{x}+\bar{y})|}{|y|^2}\le\\
&\le \frac{\nu}{25}\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}+\frac{25\pi}{\nu}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int d^3y|\bar{\omega}(\bar{x}+\bar{y})|^2)\\
&\le \frac{\nu}{25}\int d^3x|\bar{\nabla}\bar{\nabla}\bar{\omega}(\bar{x})|^2|\bar{\nabla}\bar{\omega}(\bar{x})|^{1+\varepsilon}+\frac{25\pi}{\nu}\int d^3x|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}(\int d^3y|\bar{\omega}(\bar{y})|^2).
\end{aligned}
$$

Putting all the estimates together we obtain

$$
\begin{aligned}
&\frac{1}{3+\varepsilon}\partial_t \int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon} d^3x \le \\
&\le 240\pi \int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon} d^3x \Big(\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x + 1\Big) \\
&\qquad +12\sqrt{12\pi}\int d^3x |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}\Big(\int d^3y |\bar\nabla\bar\omega(\bar x+\bar y)|^2+1\Big)+ \\
&+6.6\cdot 10^3\sqrt{3}\Big(\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x+1\Big)\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x\Big(\int|\bar\nabla\bar\omega(\bar x)|^2d^3x+1\Big)(N+1)+ \\
&\qquad +6\sqrt{12\pi}\Big(\int |\bar\nabla\bar\nabla\bar\omega(\bar x)|^2 d^3x+1\Big)\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x\Big(\int|\bar\nabla\bar\omega(\bar x)|^2d^3x+1\Big)(N+1)+ \\
&\qquad\qquad +\frac{150\sqrt{3}}{4\nu}\Big(\frac{20\pi}{3}\Big)^{\frac{5}{3}}\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x\int|\bar\nabla\bar\omega(\bar x)|^2d^3x+ \\
&\qquad\qquad\quad +\frac{150\sqrt{3}\pi}{\nu}\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x\Big(\int|\bar\omega(\bar y)|^2d^3y\Big) \\
&+(6.6\cdot 10^3\sqrt{3}+6\sqrt{12\pi})\Big(\int d^3x|\bar\nabla\bar\nabla\bar\omega(\bar x)|^2+1\Big)\Big(\int d^3x|\bar\nabla\bar\omega(\bar x)|^2+1\Big)(N+1).
\end{aligned}
\tag{35}
$$

Estimating the right hand side of (35) from above and integrating over t we obtain

$$
\begin{aligned}
&\int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon} d^3x(t_2) \le \\
&\le \Bigg(\int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon} d^3x(t_1)+ \\
&+3.55\cdot 10^4(N+1)(\max_{t\in[t_1,t_2]}(\int|\bar\nabla\bar\omega(\bar x)|^2d^3x(t)+1)(\int_{t_1}^{t_2}dt\int|\bar\nabla\bar\nabla\bar\omega(\bar x)|^2d^3x(t)+t_2-t_1)\Bigg) \\
&\cdot\exp\Bigg(\frac{2.48\cdot 10^3}{\nu}C(t_1,t_2)+ \\
&\quad +(2.28\cdot 10^3+3.55\cdot 10^4(N+1)(\max_{t\in[t_1,t_2]}\int|\bar\nabla\bar\omega(\bar x)|^2d^3x+1))\int_{t_1}^{t_2}dt\int|\bar\nabla\bar\nabla\bar\omega(\bar x)|^2d^3x+ \\
&\qquad\quad +(3.58\cdot 10^4+3.55\cdot 10^4N+\frac{3.376\cdot 10^4}{\nu})\max_{t\in[t_1,t_2]}\int|\bar\nabla\bar\omega(\bar x)|^2d^3x(t_2-t_1)+ \\
&\qquad\qquad +(3.58\cdot 10^4+3.55\cdot 10^4N)(t_2-t_1)\Bigg) \\
&\le \Big(\int |\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon} d^3x(t_1)+1\Big) \\
&\exp\Bigg(\frac{2.48\cdot 10^3}{\nu}C(t_1,t_2)+3.58\cdot 10^4(N+1)(t_2-t_1)+
\end{aligned}
\tag{36}
$$

$$+3.78\cdot 10^4(1+\int|\bar\nabla\bar\omega(\bar x)|^2d^3x(t_1))^2(N+1+\frac{1}{\nu})(t_2-t_1+\frac{1.945}{\nu})$$
$$\times\exp\left(2.45\cdot 10^5\frac{(t_2-t_1)}{\nu}+\frac{180}{\nu}C(t_1,t_2)+\frac{7.294\cdot 10^7}{\nu}e^{4.22N^{\frac{1}{2}}(t_2-t_1)}\times\right.$$
$$\left(1+\frac{3.99\cdot 10^{26}}{\nu^{4.02}}N^{2.01}C(t_1,t_2)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t_1,t_2)+1.7\cdot 10^5\frac{N}{\nu}+0.371\cdot 10^5(t_2-t_1)\right.$$
$$\left.\left.+1.066N^{\frac{1}{2}}(t_2-t_1)+|\omega|_{2+\frac{2}{3}\varepsilon}^{(2+\frac{2}{3}\varepsilon)(1-\alpha)}(t_1)\right)^5\right)\Bigg)$$

Part 5: Taking the scalar product of $|\omega(\bar x)|^{n-2}\bar\omega(\bar x)$ with the Navier-Stokes equation for ω and integrating over the whole space we obtain

$$\frac{1}{n}\partial_t\int|\omega(\bar x)|^nd^3x-\nu\int|\omega(\bar x)|^{n-2}\bar\omega(\bar x)\cdot\Delta\bar\omega(\bar x)d^3x= \tag{37}$$
$$=\frac{3}{4\pi}\int(\int_{|y|\le|\omega|_2^{-1}}+\int_{|y|>|\omega|_2^{-1}})(|\omega|^{n-2}(\bar x)\bar\omega(\bar x)\cdot\hat y)(\bar\omega(\bar x)\times\bar\omega(\bar x+\bar y)\cdot\hat y)\frac{d^3xd^3y}{|y|^3}.$$

We rewrite the absolute value of the integral over $|y|\le|\bar\omega|_2^{-1}$ on the right hand side of (37) as

$$\left|\int d^3x\int_{|y|\le|\omega|_2^{-1}}d^3y(|\omega|^{n-2}(\bar x)\bar\omega(\bar x)\cdot\hat y)(\bar\omega(\bar x)\times(\bar\omega(\bar x+\bar y)-\bar\omega(\bar x))\cdot\hat y)\frac{1}{|y|^3}\right| \tag{38}$$
$$\le 3\sqrt{6}\int d^3x\int_{|y|\le|\omega|_2^{-1}}d^3y\int_0^1ds|\bar\omega(\bar x)|^n\frac{|\bar\nabla\bar\omega(\bar x+s\bar y)|}{|y|^2}$$
$$\le 3\sqrt{6}\left(\frac{4\pi(2+\varepsilon)}{\varepsilon}\right)^{\frac{2+\varepsilon}{3+\varepsilon}}\int d^3x|\bar\omega(\bar x)|^n\int_0^1ds\left(\int_{|y|\le|\omega|_2^{-1}}d^3y|\bar\nabla\bar\omega(\bar x+s\bar y)|^{3+\varepsilon}\right)^{\frac{1}{3+\varepsilon}}$$
$$\le 3\sqrt{6}\left(\frac{4\pi(2+\varepsilon)}{\varepsilon}\right)^{\frac{2+\varepsilon}{3+\varepsilon}}\int d^3x|\bar\omega(\bar x)|^n\int_0^1\frac{ds}{s^{\frac{3}{3+\varepsilon}}}(\int_{|z|\le s|\omega|_2^{-1}}d^3z|\bar\nabla\bar\omega(\bar x+\bar z)|^{3+\varepsilon})^{\frac{1}{3+\varepsilon}}$$
$$\le 3\sqrt{6}\frac{3+\varepsilon}{\varepsilon}\left(\frac{4\pi(2+\varepsilon)}{\varepsilon}\right)^{\frac{2+\varepsilon}{3+\varepsilon}}\int d^3x|\bar\omega(\bar x)|^n(\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x)^{\frac{1}{3+\varepsilon}}$$
$$\le\frac{3\sqrt{6}(3+\varepsilon)}{\varepsilon}\left(\frac{4\pi(2+\varepsilon)}{\varepsilon}\right)^{\frac{2+\varepsilon}{3+\varepsilon}}\int|\bar\omega(\bar x)|^nd^3x(\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x)^{\frac{1}{3+\varepsilon}}$$
$$\le 4.2\cdot 10^5\int|\bar\omega(\bar x)|^nd^3x\left(\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x\right)^{\frac{1}{3+\varepsilon}}.$$

The absolute value of the integral over $|y|>|\bar\omega|_2^{-1}$ can be estimated as

$$3\sqrt{2}\int d^3x|\bar\omega(\bar x)|^n\int_{|y|>|\omega|_2^{-1}}\frac{|\bar\omega(\bar x+\bar y)|}{|y|^3}\le 3\sqrt{8\pi}N^2\int d^3x|\bar\omega(\bar x)|^n. \tag{39}$$

Therefore we obtain from (37) that

$$\frac{1}{n}\partial_t \int |\omega|^n(\bar{x}) d^3x \leq 1.1 \cdot 10^5 (N^2+1) \int |\omega|^n(\bar{x}) d^3x \ \max_{t\in[t_1,t_2]} \Big(\int |\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon} d^3x(t) + 1 \Big) \quad (40)$$

We obtain that

$$\ln \int |\omega|^n(\bar{x}) d^3x(t_2) \leq \ln \int |\omega|^n(\bar{x}) d^3x(t_1) + \quad (41)$$
$$+1.1 \cdot 10^5 n (N^2+1)(t_2-t_1) \ \max_{t\in[t_1,t_2]} \Big(\int |\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon} d^3x(t) + 1 \Big)$$

Therefore

$$\int |\omega|^n(\bar{x}) d^3x(t_2) \leq \int |\omega|^n(\bar{x}) d^3x(t_1) \quad (42)$$
$$\exp\Big(1.1 \cdot 10^5 n (N^2+1)(t_2-t_1) \max_{t\in[t_1,t_2]} \Big(|\bar{\nabla}\bar{\omega}|^{3+\varepsilon}_{3+\varepsilon}(t) + 1 \Big)\Big)$$

This implies

$$|\omega|_n(t_2) \leq |\omega|_n(t_1) \ \exp\Big(1.1 \cdot 10^5 (N^2+1)(t_2-t_1) \max_{t\in[t_1,t_2]} \Big(|\bar{\nabla}\bar{\omega}|^{3+\varepsilon}_{3+\varepsilon}(t) + 1 \Big)\Big)$$

Taking the limit $n \to \infty$ we obtain

$$|\omega|_\infty(t_2) \leq |\omega|_\infty(t_1) \ \exp\Big(1.1 \cdot 10^5 (N^2+1)(t_2-t_1) \ \max_{t\in[t_1,t_2]} \Big(|\bar{\nabla}\bar{\omega}|^{3+\varepsilon}_{3+\varepsilon}(t) + 1 \Big)\Big) \quad (43)$$
$$\leq |\omega|_\infty(t_1) \exp\Big(1.1 \cdot 10^5 (N^2+1)(t_2-t_1)(|\bar{\nabla}\bar{\omega}|^{3.01}_{3.01}(t_1) + 2) \times$$
$$\times \exp\Big(\frac{2.48 \cdot 10^3}{\nu} C(t_1,t_2) + 3.58 \cdot 10^4 (N+1)(t_2-t_1) +$$
$$+3.78 \cdot 10^4 (1 + |\bar{\nabla}\bar{\omega}|^2_2(t_1))^2 (N + 1 + \frac{1}{\nu})(t_2 - t_1 + \frac{1.945}{\nu})$$
$$\times \exp\Big(2.45 \cdot 10^5 \frac{(t_2-t_1)}{\nu} + \frac{180}{\nu} C(t_1,t_2) + \frac{7.294 \cdot 10^7}{\nu} e^{4.22 N^{\frac{1}{2}} (t_2-t_1)} \times$$
$$\times \Big(1 + \frac{3.99 \cdot 10^{26}}{\nu^{4.02}} N^{2.01} C(t_1,t_2) + \frac{3.562 \cdot 10^5}{\nu} N^{\frac{1}{2}} C(t_1,t_2) + 1.7 \cdot 10^5 \frac{N}{\nu} +$$
$$+0.371 \cdot 10^5 (t_2-t_1) + 1.066 N^{\frac{1}{2}} (t_2-t_1) + |\bar{\nabla}\bar{\omega}|^{0.76153}_{3.01}(t_1) \Big)^5 \Big)\Big)\Big).$$

As a result we obtain that if $|\omega|_2(t) \leq N$ for $t \in [0, T-\delta_N]$, then for $\delta \in (0, T-\delta_N)$

$$|\omega|_\infty(t) \leq |\omega|_\infty(\delta) \ \exp\Big(1.1 \cdot 10^5 (N^2+1) t \ \max_{t\in[\delta,T-\delta_N]} \Big(|\bar{\nabla}\bar{\omega}|^{3+\varepsilon}_{3+\varepsilon}(t) + 1 \Big)\Big) \quad (44)$$

$$
\begin{aligned}
&\leq |\omega|_\infty(\delta)\exp\Bigl(1.1\cdot 10^5(N^2+1)t(|\bar\nabla\bar\omega|_{3.01}^{3.01}(\delta)+2)\times\\
&\times\exp\Biggl(\frac{2.48\cdot 10^3}{\nu}C(t)+3.58\cdot 10^4(N+1)t+\\
&\qquad +3.78\cdot 10^4(1+|\bar\nabla\bar\omega|_2^2(\delta))^2(N+1+\frac{1}{\nu})(t+\frac{1.945}{\nu})\\
&\qquad \exp\Biggl(2.45\cdot 10^5\frac{t}{\nu}+\frac{180}{\nu}C(t)+\frac{7.294\cdot 10^7}{\nu}e^{4.22N^{\frac{1}{2}}t}\\
&\qquad\qquad \times\Biggl(1+\frac{3.99\cdot 10^{26}}{\nu^{4.02}}N^{2.01}C(t)+\frac{3.562\cdot 10^5}{\nu}N^{\frac{1}{2}}C(t)+1.7\cdot 10^5\frac{N}{\nu})+\\
&\qquad\qquad +0.371\cdot 10^5 t+1.066N^{\frac{1}{2}}t+|\bar\nabla\bar\omega|_{3.01}^{0.76153}(\delta)\Bigr)^5\Biggr)\Biggr)\Biggr).
\end{aligned}
$$

Now we shall improve the estimate on $|\omega|_\infty(t)$.

Proof of Theorem 2.

Part 1: Suppose first that there exists n_0 such that for all $n>n_0$, $|\omega|_n(t)\geq e^e$ for all $t\in[t_1,t_2]$. Let $n>n_0$.

As in the derivation of (37) we obtain

$$
\begin{aligned}
&\frac{1}{n}\partial_t\int|\omega|^n d^3x-\nu\int\bar\nabla[|\omega|^{n-2}(\bar x)\bar\omega(\bar x)](\bar\nabla\bar\omega(\bar x))d^3x \qquad (45)\\
&\qquad =\frac{3}{4\pi}\int\int[|\omega|^{n-2}(\bar x)\bar\omega(\bar x)\cdot\hat y][\bar\omega(\bar x)\times\bar\omega(\bar x+\bar y)\cdot\hat y]\frac{d^3xd^3y}{|y|^3}.
\end{aligned}
$$

Since we know from the Theorem 1 that $|\omega|_\infty(t)<\infty$ for all $t\in[t_1,t_2]$, we can estimate the right hand side of (45) as follows

$$
\begin{aligned}
&\frac{1}{n}\partial_t\int|\omega|^n d^3x\leq\frac{9\sqrt{6}}{4\pi}\int_0^1 ds\int|\omega|^n(\bar x)d^3x\int_{|y|\leq\min(|\omega|_\infty^{-1},1)}\frac{|\bar\nabla\bar\omega|(\bar x+s\bar y)}{|y|^2}d^3y\\
&+\frac{9\sqrt{2}}{4\pi}\int|\omega|^n(\bar x)d^3x\int_{1>|y|\geq\min(|\omega|_\infty^{-1},1)}\frac{|\omega|(\bar x+\bar y)}{|y|^3}d^3y+\frac{9\sqrt{2}}{4\pi}\int|\omega|^n(\bar x)d^3x\int_{|y|\geq 1}\frac{|\omega|(\bar x+\bar y)}{|y|^3}d^3y.
\end{aligned}
$$

Therefore

$$
\begin{aligned}
&\frac{1}{n}\partial_t\int|\omega|^n(\bar x)d^3x\leq 9\sqrt{6}\,\frac{3+\varepsilon}{4\pi\varepsilon}\left(\frac{4\pi(2+\varepsilon)}{\varepsilon}\right)^{\frac{2+\varepsilon}{3+\varepsilon}}\int|\omega|^n(\bar x)d^3x(\int|\bar\nabla\bar\omega(\bar x)|^{3+\varepsilon}d^3x)^{\frac{1}{3+\varepsilon}}+\\
&\qquad +9\sqrt{2}|\omega|_\infty(t)\ln(\max(|\omega|_\infty(t),1))\int|\omega|^n(\bar x)d^3x+3\sqrt{\frac{3N}{2\pi}}\int|\omega|^n(\bar x)d^3x.
\end{aligned}
$$

$$\frac{1}{n}\partial_t \int |\omega|^n(\bar{x})d^3x \leq \int |\omega|^n(\bar{x})d^3x \cdot$$
$$\cdot\left(12.78|\omega|_\infty(t)\mathrm{ln}(\max(|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}\right).$$

$$\frac{\partial_t \int |\omega|^n(\bar{x})d^3x}{\int |\omega|^n(\bar{x})d^3x} \leq n\left(12.78|\omega|_\infty(t)\mathrm{ln}(\max|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}\right).$$

$$\partial_t \mathrm{ln}\int |\omega|^n(\bar{x})d^3x \leq n\left(12.78|\omega|_\infty(t)\mathrm{ln}(\max(|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}\right).$$

$$\frac{\partial_t \mathrm{ln}\int |\omega|^n(\bar{x})d^3x}{\mathrm{ln}\int |\omega|^n(\bar{x})d^3x} \leq \frac{n\left(12.78|\omega|_\infty(t)\mathrm{ln}(\max(|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}\right)}{\mathrm{ln}\int |\omega|^n(\bar{x})d^3x}$$

$$\frac{\partial_t \mathrm{ln}\ \mathrm{ln}\int |\omega|^n(\bar{x})d^3x}{\mathrm{ln}\ \mathrm{ln}\int |\omega|^n(\bar{x})d^3x} \leq \frac{n\left(12.78|\omega|_\infty(t)\mathrm{ln}(\max(|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}\right)}{\mathrm{ln}\int |\omega|^n(\bar{x})d^3x\ \mathrm{ln}\ \mathrm{ln}\int |\omega|^n(\bar{x})d^3x}$$

$$\partial_t \mathrm{ln}\ \mathrm{ln}\ \mathrm{ln}\int |\omega|^n(\bar{x})d^3x \leq \frac{n\left(12.78|\omega|_\infty(t)\mathrm{ln}(\max(|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}\right)}{n\mathrm{ln}((\int |\omega|^n(\bar{x})d^3x)^{\frac{1}{n}})\mathrm{ln}(n\mathrm{ln}(\int |\omega|^n(\bar{x})d^3x)^{\frac{1}{n}})}$$

$$\partial_t \mathrm{ln}\ \mathrm{ln}\ \mathrm{ln}\int |\omega|^n(\bar{x})d^3x \leq \frac{12.78|\omega|_\infty(t)\mathrm{ln}(\max(|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}}{\mathrm{ln}|\omega|_n(t)(\mathrm{ln}n + \mathrm{lnln}|\omega|_n(t))}$$

Below we consider n satisfying the following condition

$$\mathrm{ln}n > \max_{t\in[t_1,t_2]}\left(12.78|\omega|_\infty(t)\mathrm{ln}(\max(|\omega|_\infty(t),1)) + 1.1\cdot 10^5|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(t) + 2.07\sqrt{N}\right), \qquad (46)$$

in fact, we shall consider even larger n, because instead of $|\omega|_\infty(t)$ we shall use the estimate (44)

$$|\omega|_\infty(t) \leq |\omega|_\infty(t_1)\exp\left(1.1\cdot 10^5(N^2+1)(t_2-t_1)\max_{t\in[t_1,t_2]}\left(|\bar{\nabla}\bar{\omega}|^{3+\varepsilon}_{3+\varepsilon}(t)+1\right)\right),$$

and instead of $|\bar{\nabla}\bar{\omega}|^{3+\varepsilon}_{3+\varepsilon}$ we shall use (36)

$$
\begin{aligned}
&\int |\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon} d^3x(t) \leq (\int |\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon} d^3x(t_1) + 1) \times \\
&\quad \times \exp \Bigg(\frac{2.48 \cdot 10^3}{\nu} C(t_1, t_2) + 3.58 \cdot 10^4 (N+1)(t_2 - t_1) + \\
&\qquad +3.78 \cdot 10^4 (1 + \int |\bar{\nabla}\bar{\omega}(\bar{x})|^2 d^3x(t_1))^2 (N + 1 + \frac{1}{\nu})(t_2 - t_1 + \frac{1.945}{\nu}) \\
&\qquad \times \exp \Bigg(\frac{2.45 \cdot 10^5}{\nu}(t_2 - t_1) + \frac{180}{\nu} C(t_1, t_2) + \frac{7.294 \cdot 10^7}{\nu^2} e^{4.22 N^{\frac{1}{2}}(t_2 - t_1)} \times \\
&\qquad\qquad \Bigg(1 + \frac{3.99 \cdot 10^{26}}{\nu^{4.02}} N^{2.01} C(t_1, t_2) + \frac{3.562 \cdot 10^5}{\nu} N^{\frac{1}{2}} C(t_1, t_2) + 1.7 \cdot 10^5 \frac{N}{\nu} + \\
&\qquad\qquad +0.371 \cdot 10^5 (t_2 - t_1) + 1.066 N^{\frac{1}{2}} (t_2 - t_1) + |\omega|^{0.76153}_{\frac{6.02}{3}}(t_1) \Bigg)^5 \Bigg) \Bigg) .
\end{aligned}
$$

Therefore consider n such that

$$
\begin{aligned}
&\ln n \geq 2.07\sqrt{N} + 12.78(|\omega|_\infty(t_1) + 1)(|\ln(|\omega|_\infty(t_1))| + 1) \times \\
&\quad \times \exp \Big(2.2 \cdot 10^5 (N^2 + 1)(t_2 - t_1 + 1)(|\bar{\nabla}\bar{\omega}(\bar{x})|^{3+\varepsilon}_{3+\varepsilon}(t_1) + 1) \\
&\qquad \times \exp \Bigg(\frac{2.48 \cdot 10^3}{\nu} C(t_1, t_2) + 3.58 \cdot 10^4 (N+1)(t_2 - t_1) + \\
&\qquad +3.78 \cdot 10^4 (1 + |\bar{\nabla}\bar{\omega}|^2_2(t_1))^2 (N + 1 + \frac{1}{\nu})(t_2 - t_1 + \frac{1.945}{\nu}) \\
&\qquad \exp \Bigg(2.45 \cdot 10^5 \frac{(t_2 - t_1)}{\nu} + \frac{180}{\nu} C(t_1, t_2) + \frac{7.294 \cdot 10^7}{\nu^2} e^{4.22 N^{\frac{1}{2}}(t_2 - t_1)} \\
&\qquad\qquad + \Bigg(1 + \frac{3.99 \cdot 10^{26}}{\nu^{4.02}} N^{2.01} C(t_1, t_2) + \frac{3.562 \cdot 10^5}{\nu} N^{\frac{1}{2}} C(t_1, t_2) + 1.7 \cdot 10^5 \frac{N}{\nu} + \\
&\qquad\qquad +0.371 \cdot 10^5 (t_2 - t_1) + 1.066 N^{\frac{1}{2}} (t_2 - t_1) + |\bar{\nabla}\bar{\omega}|^{0.76153}_{3+\varepsilon}(t_1) \Bigg)^5 \Bigg) \Bigg) \Bigg)
\end{aligned} \tag{47}
$$

For n satisfying (47) we have that

$$
\partial_t \ln \ln \ln \int |\omega|^n(\bar{x}) d^3x \leq 1. \tag{48}
$$

After integrating we obtain

$$
\ln \ln \ln \int |\omega|^n(\bar{x}, t) d^3x \leq t - t_1 + \ln \ln \ln \int |\omega|^n(\bar{x}, t_1) d^3x. \tag{49}
$$

Therefore we obtain

$$\int |\omega|^n(\bar{x},t)d^3x \le e^{e^{e^{t-t_1}}} \int |\omega|^n(\bar{x},t_1)d^3x \text{ on } t \in [t_1,t_2]. \tag{50}$$

Therefore we obtain that for all n satisfying (47)

$$|\omega|_n(t) \le e^{\frac{1}{n}e^{e^{(t-t_1)}}}|\omega|_n(t_1) \le e^{\frac{1}{n}e^{e^{(t-t_1)}}}|\omega|_\infty(t_1) \quad \text{for } t \in [t_1,t_2]. \tag{51}$$

Taking $n \to \infty$ we obtain that

$$|\omega|_\infty(t) \le |\omega|_\infty(t_1) \text{ for } t \in [t_1,t_2]. \tag{52}$$

Part 2: Suppose now that there exists $n_0 > 0$ such that for all $n > n_0$, $|\omega|_n(t)$ is a continuous function of t for all $t \in [t_1,t_2]$. Fix $n > n_0$ such that the condition (47) is satisfied.

Suppose $|\omega|_n(t_1) \ge e^e$. By continuity of $|\omega|_n(t)$ we obtain that there exists a set consisting of a union of open intervals $\cup_j I_j$ such that on each interval I_j, $|\omega|_n(t) < e^e$ for all $t \in I_j$. Let $(\cup_j I_j)^{\mathrm{c}} = \cup_k I'_k$, where each interval $I'_k = [t'_{1_k}, t'_{2_k}]$ is such that $|\omega|_n(t) \ge e^e$ for all $t \in I'_k$ and $|\omega|_n(t'_{1_k}) = e^e$ for $k > 1$ and $|\omega|_n(t'_{1_k}) = |\omega|_n(t_1)$ for $k = 1$.

Then by formula (51) we obtain on each interval I'_k

$$|\omega|_n(t) \le e^{\frac{1}{n}e^{e^{t'_{2_k}-t'_{1_k}}}} e^e \text{ for } k > 1. \tag{53}$$

For the first interval $I'_1 = [t_1, t'_{2_1}]$ we obtain that

$$|\omega|_n(t) \le e^{\frac{1}{n}e^{e^{t'_{2_1}-t_1}}} |\omega|_n(t_1). \tag{54}$$

Therefore

$$|\omega|_n(t) \le e^{\frac{1}{n}e^{e^{t_2-t_1}}} |\omega|_n(t_1) \text{ for all } t \in [t_1,t_2]. \tag{55}$$

Suppose $|\omega|_n(t_1) < e^e$. By continuity of $|\omega|_n(t)$ we obtain that there exists a set consisting of a union of open intervals $\cup_j I_j$ such that on each interval I_j, $|\omega|_n(t) < e^e$ for all $t \in I_j$. Let $(\cup_j I_j)^{\mathrm{c}} = \cup_k I'_k$ where each interval $I'_k = [t'_{1_k}, t'_{2_k}]$ is such that $|\omega|_n(t) \ge e^e$ for all $t \in [t'_{1_k}, t'_{2_k}]$ and $|\omega|_n(t'_{1_k}) = e^e$.

Then by formula (51) we obtain that on each interval I'_k

$$|\omega|_n(t) \le e^{\frac{1}{n}e^{e^{t'_{2_k}-t'_{1_k}}}} |\omega|_n(t'_{1_k}) \text{ for all } t \in [t'_{1_k}, t'_{2_k}]. \tag{56}$$

Therefore if $|\omega|_n(t_1) < e^e$ we obtain

$$|\omega|_n(t) \leq e^{\frac{1}{n}e^{e^{t_2-t_1}}} e^e \text{ for all } t \in [t_1, t_2]. \tag{57}$$

Therefore for every n satisfying the condition (47) we obtain that

$$|\omega|_n(t) \leq e^{\frac{1}{n}e^{e^{t-t_1}}} \max(|\omega|_n(t_1), e^e) \leq e^{\frac{1}{n}e^{e^{t-t_1}}} \max(|\omega|_\infty(t_1), e^e), \quad \text{for } t \in [t_1, t_2]. \tag{58}$$

Taking the limit $n \to \infty$ we obtain that

$$|\omega|_\infty(t) \leq \max(|\omega|_\infty(t_1), e^e) \text{ for all } t \in [t_1, t_2]. \tag{59}$$

Proof of Theorem 3.

Theorem 3: *Suppose* $|\omega|_\infty(0) < \infty$. *Then for any* $T > 0$ *there exists a unique solution of the 3D Navier Stokes Equation on* $[0, T]$ *such that* $|\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e) < \infty$ *for* $t \in [0, T]$ *with* $\delta = \frac{\nu}{2C|\omega|^2_\infty(0)}$, (where C is a constant independent of ω, ν, etc.)

Proof: First we shall construct an argument to show that if certain assumptions on ω hold then the solution is bounded. This argument will be used below.

Suppose we have that $|\omega|_\infty(0) < \infty$. Then by the Local Existence and Uniqueness Theorem (see Theorem 4.2 in [Ku], where we take $p = 2$ and $M_{2p} = |\omega|_\infty(0)$) there exists $t_0 = \frac{\nu}{2C|\omega|^2_\infty(0)}$ such that for $t \in [0, t_0]$, we have that there exists a unique solution such that $|\omega|_2(t) < \infty$ and also by the analyticity results in the same theorem we know that for all $\delta \in (0, t_0]$, $|\omega|_\infty(\delta) < \infty$, $|\omega|_{3+\varepsilon}(\delta) < \infty$, $|\bar{\nabla}\bar{\omega}|_2(\delta) < \infty$, $|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(\delta) < \infty$. Let $\delta = t_0$. Assume that $|\omega|_2(t) \to \infty$ as $t \to T$ and $|\omega|_2(t) < \infty$ for $t \in [0, T)$. Also assume that for all n large enough (n satisfying (47) with $t_1 = \delta$), $|\omega|_n(t)$ is continuous as a function of t for all $t \in [0, T)$. Then for any $N > N_0$ (for sufficiently large N_0) there exists $\delta_N > 0$ such that $|\omega|_2(t) \leq N$, for all $t \in [0, T - \delta_N]$. Then by Theorem 1 and Theorem 2 we obtain that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e)$ for $t \in [\delta, T - \delta_N]$. By taking $N \to \infty$ we obtain that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e)$ for $t \in [\delta, T)$. By Local Existence and Uniqueness Theorem (see Theorem 4.2 in [Ku]), we obtain that there exists $t_0 > 0$ such that there exists a unique solution on $t \in [T - \frac{t_0}{2}, T + \frac{t_0}{2}]$ that is continuous and analytic and such that $|\omega|_\infty(t) < \infty$ for $t \in [T - \frac{t_0}{2}, T + \frac{t_0}{2}]$ and then by Theorem 1 and Theorem 2 we obtain that

$$|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e) \text{ for } t \in [\delta, T + \frac{t_0}{2}]. \tag{60}$$

Thus we showed that if certain assumptions on ω hold then for any $T > 0$ the solution $\bar{\omega}(\bar{x}, t)$ is finite for all $\bar{x} \in \mathbf{R}^3$ and $t \in [0, T]$.

To show existence and uniqueness of a solution on $[0, T]$, we shall repeatedly use the Local Existence Uniqueness Theorem (Theorem 4.2 in [Ku]).

Since $|\omega|_\infty(0) < \infty$ there exits $t_0 > 0$ given by $t_0 = \frac{\nu}{2C|\omega|^2_\infty(0)}$ such that there exists a unique solution of 3D Navier Stokes Equations that is continuous and analytic on $[0, t_0]$. Let $\delta = t_0$, then $|\omega|_\infty(\delta) < \infty$, $|\bar{\nabla}\bar{\omega}|_{3+\varepsilon}(\delta) < \infty$.

Now let $t'_0 = \frac{\nu}{2C\max(|\omega|^2_\infty(\delta), e^{2e})}$ and consider $t \in [\delta, \delta + t'_0]$. Using $\bar{\omega}(\bar{x}, \delta)$ as the initial condition, we obtain that by the Local Existence and Uniqueness results there exists a unique solution which is continuous and analytic for $t \in [\delta, \delta + t'_0]$. By Theorem 1, Theorem 2 and arguments above we obtain that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e)$ for all $t \in [\delta, \delta + t'_0]$.

Repeating this argument k times, we obtain that for $t \in [\delta + (k-1)t'_0, \delta + kt'_0]$ there exists a unique solution of the 3D Navier Stokes Equation such that $|\omega|_2(t) \leq \max(|\omega|_\infty(\delta + (k-1)t'_0), e^e) \leq \max(|\omega|_\infty(\delta), e^e)$ and $|\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta + (k-1)t'_0), e^e) \leq \max(|\omega|_\infty(\delta), e^e)$.

For any $T > 0$, we can continue this argument up to k such that $k \geq \frac{T}{t'_0}$, to obtain that there exists a unique solution of the 3D Navier Stokes Equation on $[0, T]$ such that,

$$|\omega|_2(t) \leq \max(|\omega|_\infty(\delta), e^e) \text{ and } |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e) \text{ for } t \in [\delta, T]. \quad (61)$$

Proof of Theorem 4.

Theorem 4: *Suppose that* $|\omega|_4(0) < \infty$. *Then for any* $T > 0$ *there exists a unique solution of the 3D Navier Stokes Equation on* $[0, T]$.

Proof: First we construct the argument that shows that if certain assumptions on ω hold, then the solution remains bounded. This argument is used later in the proof.

Suppose we have that $|\omega|_4(0) < \infty$. Then by the Local Existence and Uniqueness Theorem (Theorem 4.2 in [Ku]) there exists $t'_0 = \frac{\nu}{2C|\omega|^2_4(0)}$ such that for $t \in [0, t'_0]$, there exists a unique solution which is continuous on $(0, t'_0]$. Therefore $|\omega|_2(t) < \infty$ for $t \in [0, t'_0]$. In addition, by analyticity results in Theorem 4.2 of [Ku] we obtain that for any $\delta \in (0, t'_0]$, $|\omega|_\infty(\delta) < \infty$, and also we have that $|\omega|_{3.01}(\delta) < \infty$ and $|\bar{\nabla}\bar{\omega}|_{3.01}(\delta) < \infty$. Suppose that $\delta = t'_0$.

Suppose that for some $T > 0$, $|\omega|_2(t) \to_{t \to T} \infty$, but $|\omega|_2(t) < \infty$ for $t \in [0, T)$. (By the argument above we know that $T > t'_0$). Suppose also that there exists n_0 (satisfying (47) with $t_1 = \delta$) such that for all $n > n_0$, $|\omega|_n(t)$ is a continuous function of t for all $t \in [0, T)$. For any $N > N_0$

(for a sufficiently large N_0), there exists $\delta_N > 0$ such that $|\omega|_2(t) \leq N$, for all $t \in [\delta, T-\delta_N]$. Then by Theorem 1, Theorem 2 and taking the limit $N \to \infty$, we obtain that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e)$ for all $t \in [\delta, T)$. By the Local Existence and Uniqueness Theorem (Theorem 4.2 in [Ku]) there exists $t_0'' > 0$ such that there exists a unique solution which is continuous and analytic on $[T - \frac{t_0''}{2}, T + \frac{t_0''}{2}]$. Thus for any $T > 0$ we obtain that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e)$ for $t \in [\delta, T]$.

To show existence and uniqueness on $[0, T]$, we shall repeatedly use the Local Existence Uniqueness Theorem. Consider $\bar{\omega}(\bar{x}, \delta)$ as the initial condition. Then there exists $t_0'' = \frac{\nu}{2C\max(|\omega|^2_\infty(\delta), e^{2e})}$ such that there exists a unique solution which is continuous and analytic on $[\delta, t_0'']$. By Theorem 1, Theorem 2 and the arguments above we have that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e)$ for $t \in [\delta, t_0'']$. Let $t \in [t_0'', 2t_0'']$. Then by the Local Existence and Uniqueness Theorem (see Theorem 4.2 in [Ku]) there exists a unique solution which is continuous and analytic on $[t_0'', 2t_0'']$. By Theorem 1, Theorem 2 and the arguments above this solution satisfies $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(t_0''), e^e) \leq \max(|\omega|_\infty(\delta), e^e)$. Repeating the argument k times we obtain that there exists a unique solution which is continuous and analytic on $[(k-1)t_0'', kt_0'']$, such that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty((k-1)t_0''), e^e) \leq \max(|\omega|_\infty(\delta), e^e)$ for $t \in [(k-1)t_0'', kt_0'']$. Thus for any $T > 0$, letting $k \geq \frac{T}{t_0''}$ we obtain that there exists a unique solution of 3D Navier Stokes Equations on $[0, T]$ such that $|\omega|_2(t) \leq |\omega|_\infty(t) \leq \max(|\omega|_\infty(\delta), e^e)$ for $t \in [0, T]$.

Bibliography

- [CF] P. Constantin, C. Fefferman, Direction of vorticity and the problem of global regularity for the Navier-Stokes equations, *IUMJ*, **Vol. 42**, (1993), 775-789.

- [FMRT] Foias, Manley, Rosa, Temam, Navier-Stokes Equations and Turbulence, *Encyclopedia of Mathematics*, Cambridge Univ Press, (2001).

- [G] Grujic, The geometric structure of the super-level sets and regularity for 3D Navier-Stokes equations, *IUMJ*, **Vol. 50**, (2001), 1309-1317.

- [GR] Grujic, Ruzmaikina, Interpolation between algebraic and geometric conditions for smoothness of the vorticity of the 3-D NSE, *IUMJ*, **Vol. 53**, **No. 4**, (2004), 1073-1080.

- [J] J. D. Jakson, Classical Electrodynamics, 2nd edition, Wiley, (1975).

- [Ku] Igor Kukavica, On the Dissipative Scale for the Navier-Stokes Equations, *IUMJ*, **Vol. 38**, **No. 3**, (1999).

- [LL] Lieb, Loss, Analysis, *Grad Studies in Math*, **14**, AMS, (1997).
- [RG] Ruzmaikina, Grujic, On depletion of the vortex-stretching term in the 3-D Navier-Stokes equations, *CMP*, **Vol. 247**, **No. 3**, (2004), 601-611.